

BROAD SPECTRUM INSECT CONTROL



TREE-äge®

(4% Emamectin Benzoate)

TREE-äge provides two-year control of listed arthropod pests in deciduous and coniferous trees and palms.

Effective for: Emerald Ash Borer; Western, Southern and Mountain Pine Beetles; Ambrosia Beetles; Gypsy Moth; Winter Moth; Flatheaded Borers; Bagworm; Leafminers (Lepidoptera, Coleoptera, Hymenoptera) and many more.

- 040-4100 1 Liter*
- 040-4105 1 Liter Case of 8

**1 Liter treats 33 trees (10" DBH)*



Why Use TREE-äge?

TREE-äge provides two-year control of Emerald Ash Borer and other labeled pests. Its unique formulation is designed to move through the xylem of trees without adversely impacting the normal cell function of the vascular tissue. TREE-äge can be used on trees grown in residential and commercial landscapes, golf courses, parks, municipal areas, and other locations that may be sensitive to application exposure.

- Two-year control of listed pests including Emerald Ash Borer, Pine Bark Beetle, and Gypsy Moth, and more
- Long residual activity backed by years of research and independent university studies
- Works with all Arborjet equipment

Provides two-year control of...



Emerald Ash Borer



Pine Bark Beetle



Gypsy Moth


...and More!

Photo Credit: EAB - David Cappaert, Bugwood.org, Pine Bark Beetle - Ron Long, Simon Fraser University, Bugwood.org

Proven Results

Independent university studies have shown TREE-äge to have high levels of efficacy and extended residual activity for two years of control against Emerald Ash Borer with a single application. In addition, it is also labeled for broad spectrum pest management.

TREE-äge Adult EAB Mortality




TREE-äge: Arborjet® Tree Injection
Imi + P: Imidacloprid + Pentra-Bark® (sprayed on the bark)
Imi: Imidacloprid (soil application)
Mauget: Mauget's Imicide® (injection caps)
Dino+ P: Dinotefuran (Safari®) + Pentra-Bark® (sprayed on the bark)
Dino: Dinotefuran (Safari®)(soil application)

Published by McCullough, D.G. et al. in USDA EAB Research & Development Review Meeting 2007

TREE-äge was the ONLY product to achieve control without loss of efficacy over time.

TREE-äge Compared to Other Treatment Methods



Caps: Imidacloprid injection caps
SD: Soil-applied dinotefuran
BSI: Bark-sprayed imidacloprid
BSD: Bark-sprayed dinotefuran
SI: Soil-applied imidacloprid
Control: Untreated

TREE-äge shows excellent 2 year control of Emerald Ash Borer compared to other treatment options and methods.

©2017 Arborjet, Inc. **Important: Always read and follow label instructions. Some crop protection products may not be registered for sale or use in all states or counties. Please check with your state or local extension service to ensure registration status.** TREE-äge® Insecticide is a Restricted Use Pesticide and must only be sold to and used by a state certified applicator or by persons under their direct supervision. TREE-äge® is a registered trademark of Arborjet, Inc. In addition to the pests noted on the TREE-äge federally registered label, Arborjet supports a FIFRA Section 2(ee) recommendation for TREE-äge insecticide to control additional bud and leaf and shoot, stem, trunk and branch pests. Please see the section 2(ee) recommendation to confirm that the recommendation is applicable in your state.