



Date: April 29, 2022  
 Product: IMA-jet, EPA Reg. No. 74578-1  
 Use: Weevils including, South American Palm Weevil (*Rhynchophorus palmarum*), Palmetto Weevil (*Rhynchophorus cruentatus fabricius*), Coconut Rhinoceros Beetle (*Oryctes rhinoceros*) and Red Palm Weevil (*Rhynchophorus ferrugineus*)  
 State/District: AL, AZ, CA, FL, GA, HI, LA, MS, NC, SC

Manufactured for Arborjet, Inc. 99 Blueberry Hill Road, Woburn, MA 01801 SCPLLABJ 1309A-L1C 1210

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

Dosage: 4.0 – 8.0 mL IMA-jet Systemic Insecticide per inch of cumulative trunk diameter at breast height (54” from the ground). Space injection holes approximately 6” apart, around the circumference of the tree.

Palms

**In palms (i.e., monocots), only one injection site is required:** locate the application site 1-3’ from the soil level and drill 4” deep into the stem.

LABEL RATES			EXAMPLE OF PALM SIZE CLASS		
Inch DBH	Milliliter /DBH”	IMA-jet per Palm	Plant Size Class	Canopy Spread	Trunk Height
5”	4	20 ml	Small	6-12’	5-19’
10”	4	40 ml	Medium	12-24’	20-30’
15”+	4	60 ml	Large	Over 24’	Over 30’

For flowering trees, make applications post bloom.

### WHEN TO TREAT

The South American palm weevil [*Rhynchophorus palmarum* (Coleoptera: Curculionidae)], larvae feed in the crown of palm trees, causing significant damage that may result in the death of meristematic tissues. If meristematic tissue is destroyed, the palm death follows on account of an inability to produce new fronds. Canary Islands date palms, *Phoenix canariensis*, is particularly susceptible to the SAPW. The palmetto weevil, *Rhynchophorus cruentatus* Fabricius is native to Florida, but is known to occur along the coast west to Texas and east to South Carolina, and until recently was considered a minor pest, attacking only severely wounded and dying trees. However, it is now known to be a pest of stressed nursery and transplanted palms as well as apparently healthy Canary Island date (*Phoenix canariensis* Hortorum ex Chabaud), Bismarck (*Bismarckia nobilis* Hildebrandt & H. Wendl.) and Latan (*Latania* spp.) palms. The Coconut Rhinoceros Beetle (*Oryctes rhinoceros*) is nonnative to Hawaii and is found to feed on Coconut palm (*Cocos nucifera*), African oil palm (*Elaeis guineensis*), date palms, as well as bananas, sugarcane, papayas, sisal, and pineapples. Best outcomes are treatments applied prior to damage to meristematic tissues (palm weevils and rhinoceros beetle and). Optimally, treatment should be made preventively at least 2 to 3 weeks before arthropods historically infest the host tree. As a result of systemic movement and longevity of IMA-jet in trees, this interval may be extended much earlier to 6 months should tree dormancy, adverse weather, management, asynchronous life cycle of pests, etc., allow earlier application timing.

IMA-jet® is a registered trademark of Arborjet, Inc. © 2022 Arborjet

2(ee) Registrant:  
 Arborjet, Inc.  
 99 Blueberry Hill Rd.  
 Woburn, MA 01801

Label Code: