

## RESTRICTED USE PESTICIDE

DUE TO ACUTE TOXICITY TO HUMANS FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# TREE-äge®

Date: September 12, 2015

Product: TREE-äge® EPA Reg. No. 100-1309-74578

Use: Additional Pests for Two-year Control in Deciduous, Coniferous, and Palm Trees

State: FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY

Manufactured for Arborjet, Inc. 99 Blueberry Hill Road, Woburn, MA 01801 SCPLLABJ 1309A-LIC 1210

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.

The user should have this recommendation in its possession at the time of use.

Use of this product according to this bulletin has not been reviewed or endorsed by the Office of the Indiana State Chemist.

This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.

### APPLICATION IN TREES

Tree Tissue	Target Pest	Application Rate <sup>1</sup>	Comments
Bud and Leaf	Cankerworms, Casebearer, Conifer Mites, Eastern Oak Looper, Elm Spanworm, Leafrollers, Linden Looper, Pine Needle Miner, Pine Tip Moth, Poplar Tentmaker, Variable Oakleaf Caterpillar, Yellownecked Caterpillar	Low to High	For best results, apply at least 2-3 weeks before the pest has historically been present. Consult with local extension agent for when this will occur in your area.
Shoot, Stem, Trunk and Branch	Ambrosia Beetles (such as Polyphagous shot hole borer; <i>Euwallacea fornicatus</i> ), <sup>2</sup> Banyan Stem Gall Wasp, Black Oak Gall Wasp, Black Turpentine Beetle, Carpenterworm, Cottonwood Twig Borer, Walnut Twig Beetle, <sup>2</sup> White Pine Weevil, Zimmerman Moth	Medium to High	For best results, apply at least 30 days before historical egg hatch or adult flight and to trees whose vascular tissue is not damaged. If vascular tissue is damaged or plugged by insect galleries, nematodes or fungi, uniform treatment and control may not be achieved.

<sup>1</sup>Use medium to high rates for remedial and longer residual control.

<sup>2</sup>Use of TREE-äge to control this insect may not result in the control of diseases vectored by the insect.

TREE-äge® is a registered trademark of Arborjet, Inc. © 2014 Arborjet

2(ee) Registrant:  
Arborjet, Inc.  
99 Blueberry Hill Rd.  
Woburn, MA 01801

Expiration Date: September 12, 2019 Label Code: